

# The $\Lambda\Lambda \rightarrow YN$ Weak Transition in Double- $\Lambda$ Hypernuclei

K. Sasaki<sup>a</sup>, T. Inoue<sup>b</sup>, and M. Oka<sup>c</sup>

<sup>a</sup>*Institute of Particle and Nuclear Studies, KEK, 1-1, Oho, Tsukuba, 305-0801  
Ibaraki, Japan*

<sup>b</sup>*Institut für Theoretische Physik, Universität Tübingen Auf der Mogenstelle 14,  
D-72076 Tübingen, Germany*

<sup>c</sup>*Department of physics, Tokyo Institute of Technology, 2-12-1, Ookayama,  
Meguro-ku, 152-8551 Tokyo, Japan*

---

## Abstract

We calculate the  $\Lambda\Lambda \rightarrow YN$  transition rate of  ${}^6_{\Lambda\Lambda}\text{He}$  by the hybrid picture, the  $\pi$  and  $K$  exchanges plus the direct quark processes. It is found that the hyperon-induced decay is weaker than the nucleon-induced decay, but the former may reveal the short-range mechanism of the weak transition and also give a clear signal of the strong  $\Delta I = 3/2$  transition. The  $\Lambda\Lambda \rightarrow YN$  transition in double- $\Lambda$  hypernucleus is complement to the  $\Lambda N \rightarrow NN$  transition as it occurs only in the  $J = 0$  channel, while the  $J = 1$  transition is dominant in the  $\Lambda N \rightarrow NN$  case.

*Key words:* Hypernuclei, Nonmesonic decay, Direct quark process

*PACS:* 21.80.+a, 12.39.-x

---

## 1 Introduction

Recent analyses of hypernuclear nonmesonic decay reveal the importance of short-range interaction of baryonic weak transition (1; 2; 3; 4; 5; 6; 7). It is, in fact, realized that the momentum transfer is large due to the  $\Lambda - N$  mass difference, and therefore the short range interaction becomes important. We proposed to treat the short range part of  $\Lambda N \rightarrow NN$  interaction using the valence quark picture of the baryon and the effective four-quark weak Hamiltonian (1; 3; 4; 7). We found that the direct quark (DQ) process gives significantly large contribution and shows qualitatively different features from the meson exchange contribution, especially in its isospin structure. The kaon exchange contribution also largely enhance the  $\Gamma_{nn}/\Gamma_{pn}$  ratio due to its spin structure (2; 4; 5; 7).

There remains, however, another problem, i.e., experimental values of the proton asymmetry from polarized hypernucleus are incompatible with the theoretical values, which is closely related to the  $J = 0$  transition amplitudes (8). The  $J = 0$  amplitude is relatively small and therefore masked by the  $J = 1$  amplitude, but it is important to clarify the breaking of the  $\Delta I = 1/2$  rule for the nonmesonic weak decay of hypernuclei. In fact, the quark model treatments of  $\Lambda N \rightarrow NN$  transition predict large breaking of the  $\Delta I = 1/2$  rule in  $J = 0$  transition channel (1; 3; 7; 9).

For the case of double- $\Lambda$  hypernuclei, the nonmesonic decay is induced not only by the nucleon but also by the other  $\Lambda$  hyperon:  $\Lambda N \rightarrow NN$  and  $\Lambda\Lambda \rightarrow YN$ . The two-body part of the nonmesonic decay rate of double- $\Lambda$  hypernuclei is classified as

$$\begin{aligned}\Gamma_{nm} &= \Gamma_{NN} + \Gamma_{YN} \\ &= \Gamma_{nn} + \Gamma_{pn}(+\Gamma_{pp}) + \Gamma_{\Lambda n} + \Gamma_{\Sigma^- p} + \Gamma_{\Sigma^0 n}\end{aligned}\quad (1)$$

where  $\Gamma_{pp}$  stands for the  $\Sigma^+ p \rightarrow pp$  transition which is available when the virtual  $\Sigma$  exists (7). The  $\Gamma_{YN}$  represents the  $\Lambda\Lambda \rightarrow YN$  transition, which is a characteristic transition for the multi- $\Lambda$  system. Experimentally, the  $\Lambda\Lambda \rightarrow YN$  transition should be distinguishable from the  $\Lambda N \rightarrow NN$  decay mode because the hyperon with high momentum is emitted once it occurs in double- $\Lambda$  hypernucleus. Thus the hyperon-induced decay mode provides us with valuable information on the  $|\Delta S| = 1$  baryon-baryon interaction (10; 11).

In sec.2, the hyperon-induced decay rate,  $\Gamma_{YN}$ , is expressed in terms of two-body amplitudes corresponding to the  $\Lambda\Lambda \rightarrow YN$  transition. In sec.3, we construct the meson exchange and DQ potential for the  $\Lambda\Lambda \rightarrow YN$  transition. In sec.4, we show the result of the  $\Lambda\Lambda \rightarrow YN$  transition rate in  ${}^6_{\Lambda\Lambda}\text{He}$ . Finally, sec.5 is devoted for the conclusion of this work.

## 2 The $\Lambda\Lambda \rightarrow YN$ transition rate

For the nonmesonic weak decay of double- $\Lambda$  hypernuclei, the hyperon induced decay mode,  $\Lambda\Lambda \rightarrow YN$ , becomes possible besides the usual  $\Lambda N \rightarrow NN$  transition. Now we concentrate on the novel  $\Lambda\Lambda \rightarrow YN$  decay. The  $\Lambda\Lambda \rightarrow YN$  transition rate is given by

$$\Gamma_{YN} = \int \frac{d^3 p'_1}{(2\pi)^3} \int \frac{d^3 p'_2}{(2\pi)^3} \frac{1}{2J_H + 1} \sum_{all} (2\pi) \delta(E_f - E_i) |M_{fi}|^2 \quad (2)$$

where  $M_{fi}$  is the  $\Lambda\Lambda \rightarrow YN$  transition amplitude,  $J_H$  is the total spin of initial hypernucleus, and  $p'_1$  and  $p'_2$  are momenta of emitted particles, *i.e.*, hyperon and nucleon. The summation indicates a sum over all quantum numbers of the initial and final particle systems. The  $\delta$ -function for the energy conservation is given by

$$\delta(E_f - E_i) = \delta \left( M_Y + M_N + \frac{p_1'^2}{2M_Y} + \frac{p_2'^2}{2M_N} - 2M_\Lambda - \epsilon_{\Lambda\Lambda} \right) \quad (3)$$

where the hyperon binding energy is taken as  $\epsilon_{\Lambda\Lambda} \simeq 7.25\text{MeV}$  (12). After the decomposition of angular momentum, the explicit form of  $M_{fi}$  is

$$|M_{fi}|^2 = (4\pi)^4 \left| \iiint \Psi_f^{L'S'J}(R, r') V_{SS'J}^{LL'}(r, r') \Psi_i^{LSJ}(R, r) r^2 dr r'^2 dr' R^2 dR \right|^2 \quad (4)$$

where  $V_{SS'J}^{LL'}(r, r')$  is the transition potential and  $\Psi^{LSJ}(R, r)$  is the wave function of the  $\Lambda\Lambda$  or  $YN$  two-body system in the configuration space. The  $L$ ,  $S$ , and  $J$  indicate the angular momentum, spin, and total spin for two-body system, respectively.

In order to extract the  $\Lambda\Lambda$  state from the double- $\Lambda$  hypernucleus, we assume, for simplicity, that the  $\Lambda\Lambda$  state with  $J = 0$  and  $I = 0$  is decoupled from the residual nuclei:

$$|_{\Lambda\Lambda}^A Z\rangle = |(\Lambda\Lambda)_{J=0}^{I=0}\rangle \otimes |^{A-2}Z\rangle \quad (5)$$

where the double- $\Lambda$  state is always in  $^1S_0$  state in the ground state. The important point of the study the  $\Lambda\Lambda \rightarrow YN$  process is that the initial  $\Lambda\Lambda$  pair is in the  $^1S_0$  state and thus the  $\Lambda\Lambda \rightarrow YN$  transition occurs only in the  $J = 0$  channel. The  $J = 0$  transition amplitude is not well studied in the  $\Lambda N \rightarrow NN$  decay, because the  $J = 1$  amplitude is much larger. In contrast, the  $\Lambda\Lambda \rightarrow YN$  decay directly probes the  $J = 0$  transition amplitude and therefore plays a complementary role.

The initial double- $\Lambda$  state is written in terms of the shell model wave function of  $0s$  state :

$$\Psi_{\Lambda\Lambda}^{000}(\vec{r}_{\Lambda_1}, \vec{r}_{\Lambda_2}) = [\phi_{0s}^b(\vec{r}_{\Lambda_1}) \times \phi_{0s}^b(\vec{r}_{\Lambda_2})]^{J=0} = [\phi_{0s}^{b_r}(\vec{r}) \times \phi_{0s}^{b_R}(\vec{R})]^{J=0} \quad (6)$$

where  $\phi_{0s}^b$  is assumed to be a Gaussian with size parameter  $b$ . The last expression of eq.(6) is the wave function in terms of the relative and center-of-mass coordinate with  $\vec{r} = \vec{r}_{\Lambda_1} - \vec{r}_{\Lambda_2}$  and  $\vec{R} = (\vec{r}_{\Lambda_1} + \vec{r}_{\Lambda_2})/2$ . The  $b_r$  and  $b_R$  are  $\sqrt{2}b$  and  $b/\sqrt{2}$ , respectively. The wave function for the final  $YN$  state is assumed

as the plane wave with  $\vec{K} = \vec{r}_Y + \vec{r}_N$  and  $\vec{k} = (m_N \vec{p}_1 - m_Y \vec{p}_2)/(m_Y + m_N)$ . The  $b$  parameter is taken as 1.9fm which fit on the r.m.s. radius of  $\Lambda$  in  ${}^6_{\Lambda\Lambda}\text{He}$  (13).

In addition, we introduce phenomenological correlation functions. For the initial  $\Lambda\Lambda$  wave function, we multiply to the uncorrelated wave function

$$f_{\Lambda\Lambda}(r) = \left(1 - e^{-\frac{r^2}{a^2}}\right)^n + br^2 e^{-\frac{r^2}{c^2}} \quad (7)$$

with  $a = 0.8\text{fm}$ ,  $b = 0.12\text{fm}^{-2}$ ,  $c = 1.28\text{fm}$ ,  $n = 1$ , according to ref.(11). On the other hand, for the final  $YN$  state, we multiply

$$f_{YN}(r) = 1 - j_0(q_c r) \quad (8)$$

where  $j_0(x)$  is a spherical Bessel function with  $q_c = 3.93 \text{ fm}^{-1}$ .

### 3 Transition potential

In order to describe the weak  $\Lambda\Lambda \rightarrow YN$  transition, we adopt the quark-meson hybrid model, which is focused on the quark degrees of freedom in a short-range region. At long and medium distances, the transition is induced mainly by the one pion exchange (OPE) and one kaon exchange (OKE) processes. The general form of  $\Lambda\Lambda \rightarrow YN$  transition potential in the meson exchange picture is given by

$$V_{\Lambda\Lambda \rightarrow YN}(\vec{q}) = g_s [\bar{u}_Y \gamma_5 u_\Lambda] \frac{1}{\vec{q}^2 + \tilde{m}_i^2} \left( \frac{\Lambda_i^2 - \tilde{m}_i^2}{\Lambda_i^2 + \vec{q}^2} \right)^2 g_w [\bar{u}_N (A + B \gamma_5) u_\Lambda] \quad (9)$$

where the coupling constants  $g_s$ ,  $g_w$ ,  $A$  and  $B$ , shown in Table 1, are chosen properly for each transition. A monopole form factor with cutoff parameters,  $\Lambda_\pi = 800 \text{ MeV}$  and  $\Lambda_K = 1300 \text{ MeV}$ , are employed for each vertex. As the energy transfer is significantly large, we introduce the effective meson mass:

$$\tilde{m} = \sqrt{m^2 - (q^0)^2}, \quad q^0 = \begin{cases} 88.5\text{MeV} & \text{for } \Lambda\Lambda \rightarrow \Lambda n \text{ transition} \\ 50.2\text{MeV} & \text{for } \Lambda\Lambda \rightarrow \Sigma N \text{ transition} \end{cases} \quad (10)$$

The kaon exchange potential can be constructed similarly. Both the strong and weak coupling constants are evaluated under the assumption of the flavor SU(3) symmetry. Note that, for OKE, it involves the strangeness transfer and thus the strong and weak vertices should be exchanged.

Table 1

The strong and weak coupling constants for the  $\pi$  and  $K$  exchange potentials. The strong coupling constants are taken from Nijmegen strong couplings (14). The weak coupling constants are in unit of  $g_w \equiv G_F m_\pi^2 = 2.21 \times 10^{-7}$ .

| Meson | Strong c.c.                     | Weak c.c.                         |                                   |
|-------|---------------------------------|-----------------------------------|-----------------------------------|
|       |                                 | PC                                | PV                                |
| $\pi$ | $g_{\Lambda\Sigma\pi} = 12.04$  | $A_{\Lambda n\pi^0} = -1.05$      | $B_{\Lambda n\pi^0} = 7.15$       |
|       |                                 | $A_{\Lambda p\pi^-} = 1.48$       | $B_{\Lambda p\pi^-} = -10.11$     |
| $K$   | $g_{\Lambda\Lambda K} = -17.65$ | $A_{\Lambda\Lambda K^0} = 0.67$   | $B_{\Lambda\Lambda K^0} = 12.72$  |
|       |                                 | $A_{\Lambda\Sigma^- K^+} = -0.55$ | $B_{\Lambda\Sigma^- K^+} = -8.42$ |
|       |                                 | $A_{\Lambda\Sigma^0 K^0} = 0.39$  | $B_{\Lambda\Sigma^0 K^0} = 5.95$  |

For shorter distances, we employ the direct quark (DQ) transition potential based on the constituent quark picture of baryons. In this mechanism, a strangeness changing weak interaction between two constituent quarks induces the transition of two baryons.

The DQ potential is given as a nonlocal form as

$$V_{DQ}^{LL'}(r, r') = -\frac{G_F}{\sqrt{2}} \times W \sum_{i=1}^7 \left\{ V_i^f f_i(r, r') + V_i^g g_i(r, r') + V_i^h h_i(r, r') \right\} \quad (11)$$

where  $r$  ( $r'$ ) stands for the radial part of the relative coordinate in the initial (final) state. The explicit forms of  $f_i$ ,  $g_i$ , and  $h_i$  are given in ref. (4), and the coefficients,  $V_i^k$ , for the  $\Lambda\Lambda \rightarrow YN$  transitions are listed in Table 2.

#### 4 Nonmesonic Decay of Double- $\Lambda$ Hypernuclei

Now we show the results of  $\Lambda\Lambda \rightarrow YN$  decay rates of  ${}^6_{\Lambda\Lambda}\text{He}$ . In Table 3, the  $\Lambda\Lambda \rightarrow YN$  transition is estimated by extending the  $\pi + K + \text{DQ}$  model to the  $S = -2$  system.

For the case of  $\Lambda\Lambda \rightarrow \Lambda n$  transition, the OPE process is forbidden because of the isospin symmetry on the strong vertex. Although the short-range interaction becomes dominant, their contributions are drastically reduced by the short range correlation. The DQ contribution is much larger than the kaon exchange contribution for the PV transition channel, but it is about 0.1% of the free  $\Lambda$  decay rate. In our full calculation, we find that the DQ reduces the total PC transition rate to almost zero. In contrast, the meson exchange

Table 2

The coefficients,  $V_i^k$ , for  $\Lambda\Lambda \rightarrow YN$  transition.

| $\Lambda\Lambda \rightarrow \Lambda n$  | $k$                 | $V_1^k$ | $V_2^k$ | $V_3^k$ | $V_4^k$ | $V_5^k$ | $V_6^k$ | $V_7^k$ |
|-----------------------------------------|---------------------|---------|---------|---------|---------|---------|---------|---------|
| $^1S_0 \rightarrow ^1S_0$               | $f$                 | -0.3142 | 0.0595  | -0.0298 | 0.0003  | -0.0298 | -0.0336 | -0.0568 |
|                                         | $\rightarrow ^3P_0$ | -0.0002 | -0.0002 | 0.0583  | 0.0212  | 0.0046  | 0.0050  | 0.0140  |
|                                         | $h$                 | -0.0002 | -0.0002 | -0.0193 | -0.0002 | -0.0143 | 0.0236  | 0.0192  |
| $\Lambda\Lambda \rightarrow \Sigma^0 n$ | $k$                 | $V_1^k$ | $V_2^k$ | $V_3^k$ | $V_4^k$ | $V_5^k$ | $V_6^k$ | $V_7^k$ |
| $^1S_0 \rightarrow ^1S_0$               | $f$                 | 0       | -0.1419 | -0.0651 | -0.0089 | 0.0282  | -0.1631 | -0.1051 |
|                                         | $\rightarrow ^3P_0$ | 0       | 0.0005  | -0.0093 | -0.0052 | 0.0083  | -0.0867 | -0.0543 |
|                                         | $h$                 | 0       | 0.0005  | 0.0035  | 0.0025  | -0.0212 | 0.1265  | 0.0800  |
| $\Lambda\Lambda \rightarrow \Sigma^- p$ | $k$                 | $V_1^k$ | $V_2^k$ | $V_3^k$ | $V_4^k$ | $V_5^k$ | $V_6^k$ | $V_7^k$ |
| $^1S_0 \rightarrow ^1S_0$               | $f$                 | 0       | 0.1004  | 0.0460  | -0.0009 | -0.0199 | 0.0720  | 0.1177  |
|                                         | $\rightarrow ^3P_0$ | 0       | 0.0011  | 0.0065  | 0.0037  | -0.0034 | 0.0391  | 0.0606  |
|                                         | $h$                 | 0       | 0.0011  | -0.0024 | 0.0043  | 0.0144  | -0.0567 | -0.0894 |

and the DQ are added coherently in the PV channels, resulting in a relatively large PV transition rate, which is about 0.2% of the free  $\Lambda$  decay.

The OPE contributions to the  $\Lambda\Lambda \rightarrow \Sigma N$  transition is expected to be small in the  $J = 0$  transition. In fact, some previous analyses have revealed that the  $J = 0$  decay rate is only few % of the  $J = 1$  transition rate. Table 3 shows that the decay rate in OPE is about 1% of the free  $\Lambda$  decay in  $\Lambda\Lambda \rightarrow \Sigma^- p$  transition. The kaon exchange process additively contributes to the OPE process, and the decay rate becomes about twice as large as the OPE alone in both the PC and PV channels. Thus the  $\Lambda\Lambda \rightarrow \Sigma N$  decay rate in meson exchange processes becomes much larger than the  $\Lambda\Lambda \rightarrow \Lambda n$ . The DQ transition rate for the  $\Lambda\Lambda \rightarrow \Sigma N$  transition is also larger than the  $\Lambda\Lambda \rightarrow \Lambda n$  case, and it behaves differently from the  $\Lambda\Lambda \rightarrow \Lambda n$  transition. In our full calculation, the PV transition rate is largely reduced, while the PC is enhanced due to the DQ contribution. Therefore the  $\Lambda\Lambda \rightarrow \Sigma N$  transition occurs mainly in the PC channel.

It is also interesting to check the breaking of the empirical  $\Delta I = 1/2$  rule for the  $\Lambda\Lambda \rightarrow YN$  transition, which leads to

$$\Gamma_{\Sigma^- p} = 2\Gamma_{\Sigma^0 n}. \quad (12)$$

This relation shows that the nonmesonic weak decay of double- $\Lambda$  hypernuclei provides us with an opportunity of direct confirmation of the validity of the empirical  $\Delta I = 1/2$  dominance. In our prediction, this relation is badly

violated due to the DQ contribution, in which the  $\Delta I = 3/2$  amplitude exists through the weak four-quark Hamiltonian. This situation is similar to the  $\Lambda N \rightarrow NN$  transition. Thus the test of the  $\Delta I = 1/2$  rule in the  $\Lambda\Lambda \rightarrow YN$  transition may reveal the importance of the short-range part in the weak baryonic interaction, and may also help to understand the mechanism of the  $\Delta I = 1/2$  enhancement observed in the nonleptonic weak decays of the free hyperons and kaons.

Table 3

$\Lambda\Lambda \rightarrow YN$  decay rates of  ${}^6_{\Lambda\Lambda}\text{He}$  in unit of  $\Gamma_\Lambda$ . The PC and PV denote the  ${}^1S_0 \rightarrow {}^1S_0$  parity-violating and  ${}^1S_0 \rightarrow {}^3P_0$  parity-conserving transitions, respectively.

| ${}^6_{\Lambda\Lambda}\text{He}$ | $\Gamma_{\Lambda n}$ |        | $\Gamma_{\Sigma^0 n}$ |        | $\Gamma_{\Sigma^- p}$ |        |
|----------------------------------|----------------------|--------|-----------------------|--------|-----------------------|--------|
|                                  | PC                   | PV     | PC                    | PV     | PC                    | PV     |
| $\pi$                            | —                    | —      | 0.0004                | 0.0030 | 0.0009                | 0.0061 |
| $\pi + K$                        | 0.0001               | 0.0002 | 0.0011                | 0.0040 | 0.0021                | 0.0079 |
| DQ                               | 0.0001               | 0.0012 | 0.0047                | 0.0037 | 0.0012                | 0.0018 |
| $\pi + K + \text{DQ}$            | 0.0000               | 0.0024 | 0.0065                | 0.0000 | 0.0064                | 0.0021 |
| total                            | 0.0024               |        | 0.0065                |        | 0.0085                |        |
| Ref.(10)                         | 0.025                |        | 0.0006                |        | 0.0012                |        |
| Ref.(11)(E)                      | 0.0044               |        | 0.011                 |        | 0.022                 |        |
| Ref.(11)(F)                      | 0.036                |        | 0.001                 |        | 0.003                 |        |

We compare our results with the previous calculations carried out by Itonaga et al. (10) and Parreño et al. (11). We all agree the bare pion-exchange amplitudes in the  $\Lambda\Lambda \rightarrow \Sigma N$  transitions (15). In Table 3, Ref. (10) includes effects of two-pion exchange in the scalar ( $\sigma$ ) channel as well as the vector ( $\rho$ ) channel. Ref. (11) contains all the pseudoscalar mesons and the vector mesons exchanged between  $\Lambda\Lambda$ . Ref.(11) took two different approaches for the final state interaction. The full calculation (F) is based on the scattering  $T$  matrix in the final states, while an effective treatment (E) evaluates the transition amplitudes by multiplying a correlation function to the relative wave function. They pointed out that the full calculation enhances  $\Gamma_{\Lambda n}$  while it reduces  $\Gamma_{\Sigma N}$ . The effective treatment does not include the conversion from  $\Sigma N$  to  $\Lambda N$ .

In the present work, we make an effective treatment of the final state interactions. Therefore, our results have to be compared to the ones in Ref.(11)(E). Our calculation is similar to their effective treatment and thus the  $\Gamma_{\Lambda n}$  enhancement due to the final state interaction is not shown. The total decay rates in our results are reduced by a factor about two due to the larger  $b$  parameter for the  $\Lambda$  wave function. We note that if the same  $b$  value is used, our results agree with the (E) results in Ref.(11). We also point out that the ratio  $\Gamma_{\Sigma^- p}/\Gamma_{\Sigma^0 n}$  is fixed to 2 by the  $\Delta I = 1/2$  rule in the meson exchange

calculations.

## 5 Conclusion

In the double  $\Lambda$  hypernuclear system, the novel decay mode,  $\Lambda\Lambda \rightarrow YN$ , is allowed. The hyperon-induced decay rates are much smaller than the decay rate of free  $\Lambda$  ( $\Gamma_\Lambda$ ). This is because the  $\Lambda\Lambda \rightarrow YN$  transition occurs only in the  $J = 0$  channel. Although the  $\Lambda\Lambda \rightarrow YN$  transition is not easy to detect experimentally, it is important to extract the information about the  $J = 0$  transition.

Our result shows that the DQ process gives significant contribution to the  $\Lambda\Lambda \rightarrow YN$  transition. It strongly suppresses the PC amplitude for the  $\Lambda\Lambda \rightarrow \Lambda n$  transition and the PV amplitude for the  $\Lambda\Lambda \rightarrow \Sigma N$  transition. Furthermore the DQ largely breaks the empirical  $\Delta I = 1/2$  rule. Unlike the  $\Lambda N \rightarrow NN$  transition, we have an opportunity to check directly the breaking of  $\Delta I = 1/2$  dominance for the  $\Lambda\Lambda \rightarrow \Sigma N$  transition.

## Acknowledgment

The authors acknowledge Profs. A. Parreño, K. Itonaga, H. Nemura, and Y. Akaishi for variable discussions. One of authors, K. S., thanks to JSPS Research Fellowship for financial support. This work is supported in part by Grants-in-Aid for Scientific Research (No.11640261) from Japan Society for the Promotion of Science (JSPS).

## References

- [1] T. Inoue, S. Takeuchi, and M. Oka, Nucl. Phys. **A577** (1994) 281c;  
*ibid* **A597** (1996) 563
- [2] A. Parreño, A. Ramos, and C. Bennhold, Phys. Rev. **C56** (1997) 339;  
A. Parreño and A. Ramos, Phys. Rev. **C65** (2002) 015204.
- [3] T. Inoue, M. Oka, T. Motoba, and K. Itonaga, Nucl. Phys. **A633** (1998) 312.
- [4] K. Sasaki, T. Inoue, and M. Oka, Nucl. Phys. **A669**, 331 (2000);  
Erratum, *ibid* **A678** (2000) 455.
- [5] D. Jido, E. Oset, and J. E. Palomar, Nucl. Phys. **A694** (2001) 525.
- [6] K. Itonaga, T. Ueda, and T. Motoba, Phys. Rev. **C65** (2002) 034617.
- [7] K. Sasaki, T. Inoue, and M. Oka, Nucl. Phys. **A707** (2002) 477.



- [8] H. Nabetani, T. Ogaito, T. Sato, and T. Kishimoto, Phys. Rev. **C60** (1999) 017001.
- [9] K. Maltman and M. Shmatikov, Phys. Lett. **B331** (1994) 1.
- [10] K. Itonaga, T. Ueda, and T. Motoba, Nucl. Phys. **A691** (2001) 197c, and private communications.
- [11] A. Parreño, A. Ramos, and C. Bennhold, Phys. Rev. **C65** (2002) 015205.
- [12] H. Takahashi et al., Phys. Rev. Lett. **87** (2001) 212502.
- [13] Y. Akaishi, and H. Nemura, private communications
- [14] Th. A. Rijken, V. G. J. Stoks, and Y. Yamamoto, Phys. Rev. **C59** (1999) 21. V. G. J. Stoks and Th. A. Rijken, Phys. Rev. **C59** (1999) 3009.
- [15] K. Itonaga, and A. Parreño, private communications